

Central Pressure: Variability and Impact of Cardiovascular Risk Factors : The Anglo-Cardiff Collaborative Trial II

Carmel M. McEniery, Yasmin, Barry McDonnell, Margaret Munnery, Sharon M. Wallace, Chloe V. Rowe, John R. Cockcroft and Ian B. Wilkinson

Hypertension. 2008;51:1476-1482; originally published online April 21, 2008;
doi: 10.1161/HYPERTENSIONAHA.107.105445

Hypertension is published by the American Heart Association, 7272 Greenville Avenue, Dallas, TX 75231
Copyright © 2008 American Heart Association, Inc. All rights reserved.
Print ISSN: 0194-911X. Online ISSN: 1524-4563

The online version of this article, along with updated information and services, is located on the World Wide Web at:

<http://hyper.ahajournals.org/content/51/6/1476>

Permissions: Requests for permissions to reproduce figures, tables, or portions of articles originally published in *Hypertension* can be obtained via RightsLink, a service of the Copyright Clearance Center, not the Editorial Office. Once the online version of the published article for which permission is being requested is located, click Request Permissions in the middle column of the Web page under Services. Further information about this process is available in the [Permissions and Rights Question and Answer](#) document.

Reprints: Information about reprints can be found online at:
<http://www.lww.com/reprints>

Subscriptions: Information about subscribing to *Hypertension* is online at:
<http://hyper.ahajournals.org/subscriptions/>

Central Pressure: Variability and Impact of Cardiovascular Risk Factors

The Anglo-Cardiff Collaborative Trial II

Carmel M. McEniery, Yasmin, Barry McDonnell, Margaret Munnery, Sharon M. Wallace, Chloe V. Rowe, John R. Cockcroft, Ian B. Wilkinson,
on Behalf of the Anglo-Cardiff Collaborative Trial Investigators

Abstract—Pulse pressure varies throughout the arterial tree, resulting in a gradient between central and peripheral pressure. Factors such as age, heart rate, and height influence this gradient. However, the relative impact of cardiovascular risk factors and atheromatous disease on central pressure and the normal variation in central pressure in healthy individuals are unclear. Seated peripheral (brachial) and central (aortic) blood pressures were assessed, and the ratio between aortic and brachial pulse pressure (pulse pressure ratio, ie, 1/amplification) was calculated in healthy individuals, diabetic subjects, patients with cardiovascular disease, and in individuals with only 1 of the following: hypertension, hypercholesterolemia, or smoking. The age range was 18 to 101 years, and data from 10 613 individuals were analyzed. Compared with healthy individuals, pulse pressure ratio was significantly increased (ie, central systolic pressure was relatively higher) in individuals with risk factors or disease ($P < 0.01$ for all of the comparisons). Although aging was associated with an increased pulse pressure ratio, there was still an average \pm SD difference between brachial and aortic systolic pressure of 11 ± 4 and 8 ± 3 mm Hg for men and women aged >80 years, respectively. Finally, stratifying individuals by brachial pressure revealed considerable overlap in aortic pressure, such that $>70\%$ of individuals with high-normal brachial pressure had similar aortic pressures as those with stage 1 hypertension. These data demonstrate that cardiovascular risk factors affect the pulse pressure ratio, and that central pressure cannot be reliably inferred from peripheral pressure. However, assessment of central pressure may improve the identification and management of patients with elevated cardiovascular risk. (*Hypertension*. 2008;51:1476-1482.)

Key Words: central pressure ■ brachial pressure ■ pulse pressure ratio ■ pulse pressure amplification ■ hypertension ■ cardiovascular risk factors

The value of brachial artery pressure as a predictor of future cardiovascular disease is firmly established.¹ Recently, greater emphasis has been placed on pulse pressure (PP), a surrogate of large artery stiffness, especially in older individuals.^{2,3} However, there is a gradual widening of PP moving from the central to the peripheral arteries,⁴ mainly because of a rise in systolic pressure. Moreover, emerging data suggest that central PP may be more closely correlated with surrogate measures of cardiovascular risk, such as left ventricular mass⁵ and carotid intima-media thickness,⁶ than brachial PP. Furthermore, central PP appears to be an independent predictor of future cardiovascular risk in selected patient groups,^{7,8} although whether it outperforms brachial PP more generally remains to be confirmed. The Conduit Artery Function Evaluation Study⁹ further highlighted the potential importance of central pressure by confirming that antihypertensive drugs can have differential effects on central and peripheral PP, which translate into differences in outcome.

The disparity between central and peripheral pressure, which is often, but perhaps misleadingly, referred to as “pressure amplification,” is driven mainly by differences in vessel stiffness and wave reflection. Previous investigations suggest that factors such as age,¹⁰ heart rate,^{11,12} and height¹³ have differential effects on central and peripheral pressure. In addition, cardiovascular risk factors such as hypercholesterolemia,¹⁴ hypertension,¹⁵ smoking¹⁶ and the metabolic syndrome¹⁷ may have greater effects on central pressure. However, the relative importance of cardiovascular risk factors and atheromatous disease, per se, on central pressure is unclear. Moreover, the variation in the gradient between central and peripheral pressure in healthy individuals has not been examined in a large unselected cohort of individuals.

Noninvasive estimation of central pressure is now possible using a number of commercial devices,¹⁸ and we have reported previously the effect of aging on the gradient

Received November 19, 2007; first decision December 7, 2007; revision accepted March 25, 2008.

From the Clinical Pharmacology Unit (C.M.M., Y., S.M.W., C.V.W., I.B.W.), University of Cambridge, Addenbrooke's Hospital, Cambridge; and the Department of Cardiology (B.M., M.M., J.R.C.), University of Cardiff, University Hospital, Cardiff, United Kingdom.

Correspondence to Carmel M. McEniery, Clinical Pharmacology Unit, University of Cambridge, Addenbrooke's Hospital Box 110, Cambridge CB2 2QQ, United Kingdom. E-mail cmm41@cam.ac.uk

© 2008 American Heart Association, Inc.

Hypertension is available at <http://hyper.ahajournals.org>

DOI: 10.1161/HYPERTENSIONAHA.107.105445

between brachial and aortic PP in a cohort of 4001 healthy individuals from the Anglo-Cardiff Collaborative Trial.¹⁰ The aim of the present study was to determine the impact of cardiovascular risk factors and cardiovascular disease on the gradient between central and peripheral PP and to investigate the variation in central pressure in healthy individuals across a wide spectrum of age and brachial pressure in a large cohort of individuals in the Anglo-Cardiff Collaborative Trial. Such information is required before studies can be designed to compare treatment strategies based on central versus peripheral pressure reduction and to define reference values for central pressure.

Methods

Subjects

Subjects were drawn from the Anglo-Cardiff Collaborative Trial population, which consists of $\approx 12\,000$ individuals selected at random from local general practice lists and open-access cardiovascular risk assessment clinics across East Anglia and Wales in the United Kingdom. The overall response rate was 85%. Subjects with secondary forms of hypertension were excluded, as were subjects in whom hemodynamic and biochemical data were incomplete at the time of analysis. This yielded a total of 11 340 individuals who were available for the current analyses. Healthy control subjects were defined as those individuals without cardiovascular disease or risk factors who were free from medication. Other subjects without overt cardiovascular disease were then grouped according to the presence of only 1 of the following risk factors: hypertension (documented or brachial systolic blood pressure [BP] ≥ 140 mm Hg and/or brachial diastolic BP ≥ 90 mm Hg), hypercholesterolemia (documented or fasting total cholesterol > 6.2 mmol/L), or smoking (> 1 cigarette per day). Subjects were excluded from these groups if > 1 risk factor was present to minimize the potentially confounding influence of individuals with multiple risk factors. Two further groups were formed: subjects with diabetes mellitus (World Health Organization criteria) but without cardiovascular disease, and those with overt cardiovascular disease (International Classification of Diseases, 10th Revision, criteria). However, because of the strong clustering of risk factors associated with each of these conditions, it was not feasible to exclude subjects with multiple risk factors. Overall, data from 10 613 individuals were included in the current analyses. Approval for all of the studies was obtained from the local research ethics committees, and written informed consent was obtained from each participant.

Hemodynamics

Seated BP was recorded in the dominant arm using a validated oscillometric technique (HEM-705CP, Omron Corporation). Radial artery waveforms were then recorded with a high-fidelity micromanometer (SPC-301, Millar Instruments) from the wrist of the dominant arm. Pulse wave analysis (SphygmoCor, AtCor Medical) was then used to generate a corresponding central (ascending aortic) waveform using a generalized transfer function,¹⁹ which has been prospectively validated for the assessment of ascending aortic BP.²⁰ Using the integral software, augmentation pressure was calculated as the difference between the second and first systolic peaks, and augmentation index was calculated as augmentation pressure expressed as a percentage of the PP. Heart rate and mean arterial pressure were obtained from the pressure waveform. All of the measurements were made in duplicate by trained investigators, and the mean values were used in the subsequent analysis. The within- and between-observer measurement reproducibility values for the augmentation index were in agreement with our previously published data.²¹

Protocol

Height and weight were assessed. After 15 minutes of seated rest, BP and radial artery waveforms were recorded. Ten milliliters of blood

were drawn from the antecubital fossa into plain tubes. The samples were centrifuged at 4°C (4000 rpm for 20 minutes), and the serum was separated and stored at -80°C for subsequent analysis. Cholesterol, triglycerides, glucose, and C-reactive protein were determined using standard methodology in an accredited laboratory.

Data Analysis

Traditionally, the term “amplification” has been used to describe the disparity between central and peripheral pressure and is calculated as peripheral PP:central PP. However, in the current study we have chosen to express aortic PP relative to brachial PP (ie, the reciprocal of PP amplification), which we refer to as the PP ratio. Thus, a higher ratio represents a relatively higher aortic pressure for a given brachial pressure. Data were analyzed using SPSS software (version 12.0). Comparisons between healthy subjects and individual risk factors were made using Student’s *t* tests with Bonferroni corrections. Analyses were performed separately for men and women. Stepwise multiple linear regression analyses were also performed to determine the factors influencing pressure amplification. Variables entered into the model were chosen if significantly associated in simple correlation analyses, and those variables known or previously associated with pressure amplification, from published observations. Values represent means \pm SDs unless otherwise stated, and a *P* value of < 0.05 was considered significant.

Results

The characteristics of the healthy subjects versus those individuals with cardiovascular risk factors or disease are presented in Table 1. With the exception of smoking, subjects with cardiovascular risk factors or disease tended to be older and heavier, have elevated brachial and aortic BPs, and have adverse biochemical profiles. In contrast, smokers were younger and had lower aortic augmentation pressure and systolic pressure than healthy subjects, although diastolic BP and triglyceride levels were higher.

Effect of Cardiovascular Risk Factors on PP Ratio

To examine the influence of cardiovascular risk factors and disease on the ratio between central and peripheral PP, comparisons were made between healthy subjects and individuals with risk factors or disease at baseline (unadjusted values) and after adjustment for confounding variables (Table 2). With the exception of smoking, all of the cardiovascular risk factors and disease were associated with an increased PP ratio (ie, aortic pressure was relatively higher in these individuals) compared with healthy individuals at baseline. However, after adjustment for differences in age, height, and heart rate between the groups, all of the risk factors and cardiovascular disease, per se, were associated with an increased PP ratio in men and women ($P < 0.01$ for all of the comparisons). Diabetes and cardiovascular disease in men and hypertension, diabetes, and cardiovascular disease in women were associated with the most marked elevation in the PP ratio. In addition, as a posthoc analysis, the influence of obesity on the PP ratio was examined by comparing healthy individuals with a BMI of < 25 kg/m² with those with a BMI of > 30 kg/m² and without additional risk factors. After adjustment for confounding variables, the PP ratio did not differ between nonobese and obese men (0.70 ± 0.10 versus 0.70 ± 0.08 , nonobese versus obese; $P = 0.5$) but was significantly increased in obese women (0.72 ± 0.10 versus 0.74 ± 0.09 , nonobese versus obese; $P = 0.003$).

Table 1. Characteristics of Subjects Grouped According to the Presence of Cardiovascular Risk Factors or Disease

Parameter	Healthy (n=5648)	Hypertension (n=3420)*	Hypercholesterolemia (n=289)*	Smoking (n=290)*	Diabetes (n=356)	Cardiovascular Disease (n=610)
Age, y	45±21	61±17‡	57±16‡	33±16‡	65±14‡	70±10‡
Age range, y	18 to 92	18 to 94	18 to 81	18 to 101	18 to 85	38 to 92
Gender, male/female	2779/2869	1990/1430	117/172	153/137	128/228	167/443
Height, m	1.69±0.10	1.69±0.09	1.67±0.09‡	1.71±0.10‡	1.68±0.09‡	1.69±0.08
Weight, kg	72±14	78±15‡	72±14	73±14	83±16‡	79±14‡
Brachial SBP, mm Hg	120±11	153±17‡	124±10‡	120±11	143±21‡	142±23‡
Brachial DBP, mm Hg	74±8	87±11‡	77±7‡	75±7‡	79±11‡	78±13‡
Brachial PP, mm Hg	46±10	66±17‡	47±10	45±10‡	63±20‡	64±20‡
Aortic SBP, mm Hg	108±12	140±17‡	115±11‡	106±10‡	130±21‡	130±23‡
Aortic DBP, mm Hg	75±8	88±11‡	78±8‡	76±7‡	80±12‡	79±13‡
Aortic PP, mm Hg	33±10	52±17‡	37±9‡	30±8‡	50±19‡	52±19‡
Aortic:brachial PP (ratio)	0.72±0.12	0.79±0.11‡	0.77±0.10‡	0.66±0.10‡	0.78±0.10‡	0.80±0.09‡
Brachial-aortic SBP, mm Hg	12±6	12±7‡	10±5‡	14±6‡	13±6‡	12±5
PP amplification (ratio)	1.44±0.25	1.30±0.20‡	1.32±0.18‡	1.54±0.22‡	1.30±0.18‡	1.28±0.16‡
MAP, mm Hg	90±8	110±11‡	93±8‡	90±8	101±14‡	100±15‡
Heart rate, bpm	69±12	70±12‡	70±10	73±12‡	71±14‡	67±14‡
Augmented pressure, mm Hg	7±7	16±10‡	10±6‡	4±5‡	14±9‡	16±10‡
Augmentation index, %	18±18	28±14‡	26±13‡	11±16‡	26±12‡	29±11‡
Total cholesterol, mmol/L	4.32±0.89	4.90±0.79‡	6.38±1.05‡	4.34±0.88	4.66±1.05‡	4.61±1.09‡
LDL, mmol/L	2.41±0.75	2.85±0.72‡	4.09±1.01‡	2.40±0.77	2.58±0.85‡	2.61±0.93‡
HDL, mmol/L	1.43±0.38	1.36±0.40‡	1.50±0.46‡	1.39±0.40	1.25±0.41‡	1.28±0.40‡
Triglycerides, mmol/L	1.13±0.70	1.58±0.94‡	1.80±0.98‡	1.30±0.81‡	2.01±1.30‡	1.69±1.02‡
Glucose, mmol/L	4.93±0.90	5.44±1.07‡	5.37±1.22‡	4.93±0.79	9.03±4.22‡	6.13±2.33‡
CRP, mg/L	2.23±4.60	2.76±4.93‡	1.91±2.46	2.33±3.58	3.54±7.47‡	5.50±28.20‡

Data are means±SDs unless otherwise specified. SBP indicates systolic blood pressure; DBP, diastolic blood pressure; PP, pulse pressure; MAP, mean arterial pressure; PP amplification; brachial PP:aortic PP; LDL, low-density lipoprotein; HDL, high-density lipoprotein; CRP, C-reactive protein.

*Data show subjects in whom hypertension, hypercholesterolemia, or smoking was the only cardiovascular risk factor: individuals in these categories with >1 risk factor were excluded from the analyses.

† $P<0.01$ versus healthy subjects.

‡ $P<0.001$ versus healthy subjects.

Factors Influencing PP Ratio

Stepwise multiple regression models were then constructed to determine those factors that independently influence the PP ratio and also the difference between brachial and aortic systolic pressures (Table 3). In addition to age, heart rate, gender, and height, all of the cardiovascular risk factors and the presence of cardiovascular disease remained independently associated with PP ratio (adjusted $R^2=0.73$; $P<0.001$) and the difference between brachial and aortic systolic pressure (adjusted $R^2=0.44$; $P<0.001$).

Interindividual Variability in PP Ratio

With aging, the PP ratio significantly increased, indicating that there was a greater age-related rise in aortic PP relative to brachial PP. Therefore, the absolute difference between aortic and brachial systolic pressures declined significantly with age in both men and women ($P<0.001$ for all comparisons; Figure 1). Despite this, there was still significant disparity between aortic and brachial PP in both men and women in the oldest age category. Indeed, there was an average difference in brachial and aortic SBP of 11 ± 5

8 ± 3 mm Hg for men and women, respectively. Moreover, within all of the age groups, there was substantial interindividual variation in the PP ratio for all of the age categories, leading to considerable overlap between age categories. For example, in terms of PP ratio, 41% of men and 28% of women aged <20 years had values that overlapped with men and women aged 40 to 49 years, respectively. Similarly, 48% of men and 46% of women aged 40 to 49 years had values that overlapped with men and women aged 70 to 79 years, respectively.

Variation in Aortic Systolic Pressure With Increasing Levels of Brachial Systolic Pressure

Given the substantial variability in the PP ratio, we investigated the interindividual variability in aortic systolic pressure as a function of brachial systolic pressure. Categories were based on 10-mm Hg increments of brachial systolic BP within the 2007 European Society of Hypertension and of the European Society of Cardiology definitions of BP²² (Figure 2). Throughout the range of categories for brachial systolic pressure, there was substantial interindividual variation in

Table 2. Influence of Cardiovascular Risk Factors and Disease on PP Ratio

Model	Men		Women	
	Data, Mean±SD	P	Data, Mean±SD	P
Healthy				
Unadjusted	0.70±0.12		0.73±0.13	
Fully adjusted	0.72±0.10		0.76±0.11	
Hypertensive				
Unadjusted	0.76±0.11	<0.001	0.83±0.09	<0.001
Fully adjusted	0.74±0.09	<0.001	0.81±0.06	<0.001
Hypercholesterolemia				
Unadjusted	0.76±0.10	<0.001	0.78±0.10	<0.001
Fully adjusted	0.74±0.06	0.001	0.77±0.09	<0.01
Smoking				
Unadjusted	0.64±0.09	<0.001	0.68±0.11	<0.001
Fully adjusted	0.74±0.10	0.001	0.78±0.08	<0.001
Diabetes				
Unadjusted	0.76±0.10	<0.001	0.80±0.10	<0.001
Fully adjusted	0.75±0.07	<0.001	0.80±0.05	<0.001
Cardiovascular disease				
Unadjusted	0.78±0.09	<0.001	0.82±0.09	<0.001
Fully adjusted	0.76±0.06	<0.001	0.80±0.06	<0.001

Fully adjusted refers to data adjusted for age, height, and heart rate. *P* values refer to comparisons with healthy subjects.

aortic systolic pressure in both men and women. Again, this variability led to extensive overlap in aortic systolic pressures between categories, without any overlap between brachial pressures, which was more pronounced in men than in women. For example, the overlap in aortic systolic BP between subjects with normal BP and those with high-normal BP was 78% in men and 63% in women. Similarly, the overlap in aortic systolic BP between high-normal BP and stage 1 hypertension was 78% in men and 73% in women. The overlap in aortic systolic BP between individuals with normal BP and stage 1 hypertension was 32% for men and 10% for women.

Discussion

The aim of the current study was to assess the extent to which the disparity between central and peripheral pressure is affected by cardiovascular risk factors and cardiovascular disease, per se, and to examine the degree of variation in central pressure in healthy individuals. Our major, novel findings were that individual cardiovascular risk factors and atherosclerosis were all associated with an increased PP ratio in both men and women, and all of these factors remained independently associated with PP ratio in multiple regression analyses. In addition, although PP ratio increased with increasing age, differences between central and peripheral PP were apparent, even in the oldest individuals, and there was also considerable interindividual variability between individuals. Finally, stratifying individuals by brachial BP highlighted the considerable overlap in aortic systolic pressure between discrete brachial BP categories, such that >70% of individuals with high-normal brachial systolic BP had similar

aortic systolic pressures to those individuals with stage 1 hypertension. Overall, these data show that the PP ratio depends on a number of factors, and that there is significant variation in central pressure between individuals despite similar brachial pressures. This suggests that central pressure cannot be reliably inferred from measurements of brachial pressure, and that measuring central pressure may improve the identification and management of patients with elevated cardiovascular risk.

The disparity between central and peripheral PP is mainly determined by differences in vessel stiffness and wave reflection. Therefore, any factor increasing wave reflections is likely to increase central pressure independently of brachial pressure. Indeed, several studies have suggested that cardiovascular risk factors such as hypercholesterolemia,¹⁴ hypertension,¹⁵ smoking,¹⁶ and the metabolic syndrome¹⁷ may have a greater effect on central pressure. However, these previous studies did not control for the inclusion of patients with multiple cardiovascular risk factors, or they examined the impact of various physiological factors, such as heart rate, on the PP gradient within risk-factor groups. Therefore, the relative impact of different cardiovascular risk factors or disease on central pressure is unclear. In the current study, all of the cardiovascular risk factors and the presence of cardiovascular disease were associated with a significantly increased PP ratio in both men and women compared with healthy individuals, even after adjusting for the potentially confounding effects of differences in age, height, and heart rate between the groups. Diabetes and cardiovascular disease in men and hypertension, diabetes, and cardiovascular disease in women were associated with more profound increases in

Table 3. Stepwise Regression Analyses

Model Regression Coefficient		SE	β	P	R ² Change, %
Aortic:brachial PP ratio (adjusted R ² =0.73, P<0.001)					
Age	0.007	<0.001	0.636	<0.001	54
Heart rate	-0.007	<0.001	-0.374	<0.001	9
Gender	0.087	0.003	0.181	<0.001	3
Height	-0.277	0.019	-0.112	<0.001	2
Hypertension	0.034	0.003	0.069	<0.001	1
Cardiovascular disease	0.051	0.006	0.051	<0.001	1
Smoking	0.032	0.007	0.025	<0.001	1
Hypercholesterolemia	0.018	0.006	0.019	0.001	1
Diabetes	0.017	0.007	0.013	0.022	1
Brachial-aortic systolic pressure difference (adjusted R ² =0.44; P<0.001)					
Age	-0.124	0.003	-0.432	<0.001	19
Gender	-3.151	0.118	-0.257	<0.001	9
Heart rate	0.168	0.004	0.330	<0.001	8
Hypertension	-1.854	0.685	-0.135	<0.001	2
Height	8.015	1.824	0.134	<0.001	2
Cardiovascular disease	-0.900	0.223	-0.036	0.001	1
Smoking	-0.845	0.254	-0.026	0.001	1
Diabetes	-0.863	0.265	-0.026	0.001	1
Hypercholesterolemia	-0.419	0.210	-0.017	0.046	1

the PP ratio than the other risk factors. In addition, the PP ratio was increased in obese subjects without any additional risk factors. Overall, these data suggest that, for a given level of brachial BP, central (aortic) BP is higher in those individuals with risk factors or disease compared with healthy subjects.

Other factors that affect the disparity between central and peripheral PP have been identified, particularly age, heart rate, and height.^{10-13,23} Consistent with these previous data, we observed that the major, independent influences on PP ratio in the current study were age, heart rate, height, and gender. In addition, all of the cardiovascular risk factors and the presence of cardiovascular disease remained independently associated with the PP ratio after adjusting for confounding variables. Our data are in close agreement with those of a previous study in which stepwise multiple regres-

sion analyses were used to determine the most important influences on PP amplification (brachial:aortic PP).²³ However, in the current study and in that of Camacho et al,²³ only ≈70% of the variability in the PP ratio was explained by the regression models. Given that a 2- to 3-mm Hg difference in brachial BP results in a 20% to 30% difference in cardiovascular risk,²⁴ these data suggest that central BP cannot be predicted with sufficient accuracy from brachial pressure by a statistical model, but rather, needs to be assessed directly, using appropriate methodologies.

Aging exerts a powerful influence on PP gradient,^{10,25} and we have confirmed this finding in the current study. We have also demonstrated that aging exerts a marked influence on the absolute difference between brachial and aortic systolic pressure in healthy subjects. However, despite this age-related effect, differences between brachial and aortic pres-




Figure 1. The effect of age on PP ratio (left) and the difference between brachial and aortic SBP (right) for healthy men (■; n=2779) and women (●; n=2869; total n=5648). The data represent means±2 SD, thus, ≈95% of the data lie within these limits.




Figure 2. Box plot of aortic systolic pressure per 10-mm Hg increments in brachial systolic pressure in men (left, $n=3603$) and women (right, $n=3176$). Analyses were performed in all of the healthy subjects and treatment-naïve hypertensive subjects. The vertical line within the box represents the median, the box represents the interquartile range (50% of the distribution), and the whiskers represent the range of values. The dashed lines indicate BP classifications according to the 2007 European Society of Hypertension and of the European Society of Cardiology guidelines.²²

tures of ≈ 11 mm Hg for men and ≈ 8 mm Hg for women were still evident in healthy individuals over the age of 80 years, disproving claims made previously that the central-to-peripheral pressure gradient is negligible in older individuals.²⁶ Moreover, there was substantial variation in the PP ratio between individuals of a similar age, which remained consistent across the entire age spectrum. These data further support previous observations that peripheral and central pressures are not the same, even in older individuals.

The marked variation in the PP ratio between individuals is highlighted in Figure 2, where there is considerable overlap in aortic systolic pressures between discrete categories of brachial systolic BP. This variability in aortic pressure for given levels of brachial pressure is likely to be meaningful clinically, because current data suggest that central, rather than brachial, pressure correlates more closely with surrogate markers of cardiovascular risk^{5,6} and independently predicts future cardiovascular events.^{7,8} There are also important implications for the categorization of hypertension, because if central BP is more important in defining an individual's risk and/or the impact of therapy, then categories that are based on central, rather than peripheral, pressure may be more useful. For example, according to current definitions for the classification of hypertension based on brachial BP,²² $>78\%$ of men with normal brachial BP had aortic systolic BP in common with men classified as having high-normal brachial BP (63% of women), and, again, 78% of men with high-normal brachial BP had aortic systolic BP in common with men with Stage 1 hypertension (73% of women). Moreover, 32% of men and 10% of women with normal brachial BP had aortic systolic BP in common with individuals with stage 1 hypertension. These observations suggest that a large proportion of individuals who are classified as being normotensive based on current guidelines might actually be at increased risk according to their central BP. Conversely, some individuals labeled as being hypertensive or at increased risk of developing hypertension might actually have lower cardiovascular risk, because they have a lower central BP.

A limitation of the current study is its cross-sectional design, and further longitudinal follow-up studies are re-

quired to confirm the age-related changes in the PP ratio described here. In addition, many of the patients with hypertension, diabetes, or overt cardiovascular disease were taking vasoactive medication, which may have influenced the degree of disparity between central and peripheral pressure observed in these subjects. However, none of the healthy subjects, hypercholesterolemic subjects, or smokers were taking vasoactive medication. Central (aortic) BP was derived noninvasively, using a generalized transfer function, which has been criticized in the past.^{27,28} However, the transfer function has been prospectively validated and is accurate to within 1.0 mm Hg,^{20,29} and our previous data demonstrate a tight correlation between derived aortic and measured carotid indices.¹⁰ Moreover, our key findings focus on the ratio between central and peripheral PP. As a fiducial parameter, this is independent of any systematic calibration errors inherent in the brachial cuff pressure measurement, relying instead on the information contained within the arterial pressure waveform.

Perspectives

Current guidelines for the diagnosis and treatment of hypertension are based solely on brachial BP. However, brachial and central BPs are not the same, even in older individuals, as demonstrated by the current data where differences of 8 to 10 mm Hg between brachial and aortic systolic BP are standard. Increased central BP is associated with a number of pathophysiological mechanisms, such as left ventricular hypertrophy,⁵ altered myocardial perfusion,³⁰ and carotid artery remodeling,⁶ all of which increase the risk of cardiovascular events. Moreover, central pressure may be a better predictor of future cardiovascular risk in selected patient groups than brachial pressure.^{7,8} Therefore, it seems likely that the assessment of central pressure will improve the identification and management of patients with elevated cardiovascular risk. However, if central pressure is ever to replace brachial pressure in clinical decision-making, new guidelines that focus on central pressure will be required. In addition, the acceptance and, ultimately, success of such guidelines will greatly depend on the widespread use of devices to assess

central pressure in primary care settings, a situation that may be considered somewhat unlikely, at least in the short-term. Therefore, further studies are urgently required to confirm the current data and to provide evidence that treatment decisions based on measurements of central BP result in better outcomes.

Acknowledgments

The Anglo-Cardiff Collaborative Trial Study Investigators include John Cockcroft, Zahid Dhakam, Stacey Hickson, Kaisa Maki-Petaja, Barry McDonnell, Carmel McEniery, Maggie Munnerly, Pawan Pusalkar, Christopher Retallick, Chloe Rowe, James Sharman, Rachel Stainsby, Edna Thomas, Sharon Wallace, Ian Wilkinson, Susannah Williams, and Yasmin.

Sources of Funding

C.M.M. is supported by a British Heart Foundation Intermediate Research Fellowship, and I.B.W. is supported by a British Heart Foundation Senior Clinical Fellowship. The Clinical Pharmacology Unit is supported by a Department of Health Biomedical Research Centre grant.

Disclosures

None.

References

- MacMahon S, Peto R, Cutler J, Collins R, Sorlie P, Neaton J, Abbott R, Godwin J, Dyer A, Stamler J. Blood pressure, stroke and coronary heart disease. Part 1, prolonged differences in blood pressure: prospective observational studies corrected for regression dilution bias. *Lancet*. 1990; 335:765–774.
- Franklin SS, Khan SA, Wong ND, Larson MG, Levy D. Is pulse pressure useful in predicting risk for coronary heart disease? The Framingham Heart Study. *Circulation*. 1999;100:354–360.
- Wilkinson IB, Franklin SS, Cockcroft JR. Nitric oxide and the regulation of large artery stiffness: from physiology to pharmacology. *Hypertension*. 2004;44:112–116.
- Kroeker EJ, Wood EH. Comparison of simultaneously recorded central and peripheral arterial pressure pulses during rest, exercise and tilted position in man. *Circ Res*. 1955;3:623–632.
- Covic A, Goldsmith DJA, Panaghiu L, Covic M, Sedor J. Analysis of the effect of haemodialysis on peripheral and central arterial pressure waveforms. *Kidney Int*. 2000;57:2634–2643.
- Boutouyrie P, Bussy C, Lacolley P, Girerd X, Laloux B, Laurent S. Association between local pulse pressure, mean blood pressure, and large-artery remodeling. *Circulation*. 1999;100:1387–1393.
- Safar ME, Blacher J, Pannier B, Guerin AP, Marchais SJ, Guyonvarc'h PM, London GM. Central pulse pressure and mortality in end-stage renal disease. *Hypertension*. 2002;39:735–738.
- Roman MJ, Devereux RB, Kizer JR, Lee ET, Galloway JM, Ali T, Umans JG, Howard BV. Central pressure more strongly relates to vascular disease and outcome than does brachial pressure: the Strong Heart Study. *Hypertension*. 2007;50:197–203.
- Williams B, Lacy PS, Thom SM, Cruickshank K, Stanton A, Collier D, Hughes AD, Thurston H, O'Rourke M. Differential impact of blood pressure-lowering drugs on central aortic pressure and clinical outcomes: principal results of the Conduit Artery Function Evaluation (CAFE) study. *Circulation*. 2006;113:1213–1225.
- McEniery CM, Yasmin, Hall IR, Qasem A, Wilkinson IB, Cockcroft JR. Normal vascular aging: differential effects on wave reflection and aortic pulse wave velocity: the Anglo-Cardiff Collaborative Trial (ACCT). *J Am Coll Cardiol*. 2005;46:1753–1760.
- Albaladejo P, Copie X, Boutouyrie P, Laloux B, Declere AD, Smulyan H, Benetos A. Heart rate, arterial stiffness, and wave reflections in paced patients. *Hypertension*. 2001;38:949–952.
- Wilkinson IB, Mohammad NH, Tyrrell S, Hall IR, Webb DJ, Paul VE, Levy T, Cockcroft JR. Heart rate dependency of pulse pressure amplification and arterial stiffness. *Am J Hypertens*. 2002;15:24–30.
- London GM, Guerin AP, Pannier BM, Marchais SJ, Metivier F. Body height as a determinant of carotid pulse contour in humans. *J Hypertens*. 1992;10:93–95.
- Wilkinson IB, Prasad K, Hall IR, Thomas A, MacCallum H, Webb DJ, Frenneaux MP, Cockcroft JR. Increased central pulse pressure and augmentation index in subjects with hypercholesterolemia. *J Am Coll Cardiol*. 2002;39:1005–1011.
- Wilkinson IB, Franklin SS, Hall IR, Tyrrell S, Cockcroft JR. Pressure amplification explains why pulse pressure is unrelated to risk in young subjects. *Hypertension*. 2001;38:1461–1466.
- Mahmud A, Feely J. Effect of smoking on arterial stiffness and pulse pressure amplification. *Hypertension*. 2003;41:183–187.
- Protogerou AD, Blacher J, Mavrikakis M, Lekakis J, Safar ME. Increased pulse pressure amplification in treated hypertensive subjects with metabolic syndrome. *Am J Hypertens*. 2007;20:127–133.
- Laurent S, Cockcroft JR, van Bortel LM, Boutouyrie P, Giannattasio C, Hayoz D, Pannier B, Vlachopoulos C, Wilkinson IB, Struijker Boudier HA. Abridged version of the expert consensus document. *Artery Res*. 2007;1:2–12.
- Karamanoglu M, O'Rourke MF, Avolio AP, Kelly RP. An analysis of the relationship between central aortic and peripheral upper limb pressure waves in man. *Eur Heart J*. 1993;14:160–167.
- Pauca AL, O'Rourke MF, Kon ND. Prospective evaluation of a method for estimating ascending aortic pressure from the radial artery pressure waveform. *Hypertension*. 2001;38:932–937.
- Wilkinson IB, Fuchs SA, Jansen IM, Spratt JC, Murray GD, Cockcroft JR, Webb DJ. Reproducibility of pulse wave velocity and augmentation index measured by pulse wave analysis. *J Hypertens*. 1998;16: 2079–2084.
- Mansia G, De Backer G, Dominiczak A, Cifkova R, Fagard R, Germano G, Grassi G, Heagerty AM, Kjeldsen SE, Laurent S, Narkiewicz K, Ruilope L, Rynkiewicz A, Schmieder RE, Struijker Boudier HA, Zanchetti A. 2007 ESH-ESC Guidelines for the management of arterial hypertension: the task force for the management of arterial hypertension of the European Society of Hypertension (ESH) and of the European Society of Cardiology (ESC). *Blood Press*. 2007;16:135–232.
- Camacho F, Avolio A, Lovell NH. Estimation of pressure pulse amplification between aorta and brachial artery using stepwise multiple regression models. *Physiol Meas*. 2004;25:879–889.
- Lewington S, Clarke R, Qizilbash N, Peto R, Collins R. Age-specific relevance of usual blood pressure to vascular mortality: a meta-analysis of individual data for one million adults in 61 prospective studies. *Lancet*. 2002;360:1903–1913.
- Sharman JE, McEniery CM, Dhakam ZR, Coombes JS, Wilkinson IB, Cockcroft JR. Pulse pressure amplification during exercise is significantly reduced with age and hypercholesterolemia. *J Hypertens*. 2007;25: 1249–1254.
- Dart AM, Cameron JD, Gatzka CD, Willson K, Liang YL, Berry KL, Wing LM, Reid CM, Ryan P, Beilin LJ, Jennings GL, Johnston CI, McNeil JJ, Macdonald GJ, Morgan TO, West MJ, Kingwell BA. Similar effects of treatment on central and brachial blood pressures in older hypertensive subjects in the Second Australian National Blood Pressure Trial. *Hypertension*. 2007;49:1242–1247.
- Davies JI, Band MM, Pringle S, Ogston S, Struthers AD. Peripheral blood pressure measurement is as good as applanation tonometry at predicting ascending aortic blood pressure. *J Hypertens*. 2003;21:571–576.
- Cloud GC, Rajkumar C, Kooner J, Cooke J, Bulpitt CJ. Estimation of central aortic pressure by SphygmoCor requires intra-arterial peripheral pressures. *Clin Sci (Lond)*. 2003;105:219–225.
- Sharman JE, Lim R, Qasem AM, Coombes JS, Burgess MI, Franco J, Garrahy P, Wilkinson IB, Marwick TH. Validation of a generalized transfer function to noninvasively derived central blood pressure during exercise. *Hypertension*. 2006;47:1203–1208.
- Recchia FA, Senzaki H, Saeki A, Byrne BJ, Kass DA. Pulse pressure-related changes in coronary flow in vivo are modulated by nitric oxide and adenosine. *Circ Res*. 1996;79:849–856.